


HOW TO USE THE KINERET® (anakinra) INJECTION TRACKER

KINERET® (anakinra) INJECTION TRACKER

Print a new injection tracker sheet every time you open a new shipment of Kineret injections.



Note the injection site each day in the log below

WEEK 1		WEEK 2		WEEK 3		WEEK 4	
DATE	8/1	DATE	8/8	DATE	8/15	DATE	
DAY 1	SITE A	DAY 8	SITE L	DAY 15	SITE B	DAY 22	SITE
DAY 2	SITE G	DAY 9	SITE C	DAY 16	SITE M	DAY 23	SITE
DAY 3	SITE M	DAY 10	SITE F	DAY 17	SITE G	DAY 24	SITE
DAY 4	SITE E	DAY 11	SITE D	DAY 18	SITE	DAY 25	SITE
DAY 5	SITE B	DAY 12	SITE A	DAY 19	SITE	DAY 26	SITE
DAY 6	SITE H	DAY 13	SITE K	DAY 20	SITE	DAY 27	SITE
DAY 7	SITE J	DAY 14	SITE H	DAY 21	SITE	DAY 28	SITE

Still have questions about injecting Kineret or injection site reactions?

Contact KINERET® On TRACK™, read the Kineret Prescribing Information, and/or ask a healthcare professional.

CALL US: 866.547.0644, 8AM to 8PM EST, Monday through Friday.



This tool has been developed to help you keep track of your child's injection sites.



Every time you open a new shipment of Kineret injections, print out an injection tracker sheet.



- Every week, note the date.
- Every day, make a note of the site where Kineret is injected.



Ask for help if it is needed!

- Injection site reactions are expected, but there are ways to address them.
- Call us: 866.547.0644, 8AM to 8PM EST, Monday through Friday.


Injection site rotation is an important factor for success with Kineret.

- Be sure to choose a new site each time Kineret is administered.
 - Choosing a new site may help avoid soreness at one particular site.
- Do not inject into an area that is tender, red, bruised or hard.
- Avoid areas with scars or stretch marks.
- Do not inject close to a vein that can be seen under the surface of the skin.



KINERET® (anakinra) INJECTION TRACKER

Print a new injection tracker sheet every time you open a new shipment of Kineret injections.



Note the injection site each day in the log below

WEEK 1		WEEK 2		WEEK 3		WEEK 4	
DATE	DATE	DATE	DATE	DATE	DATE	DATE	DATE
DAY 1	SITE	DAY 8	SITE	DAY 15	SITE	DAY 22	SITE
DAY 2	SITE	DAY 9	SITE	DAY 16	SITE	DAY 23	SITE
DAY 3	SITE	DAY 10	SITE	DAY 17	SITE	DAY 24	SITE
DAY 4	SITE	DAY 11	SITE	DAY 18	SITE	DAY 25	SITE
DAY 5	SITE	DAY 12	SITE	DAY 19	SITE	DAY 26	SITE
DAY 6	SITE	DAY 13	SITE	DAY 20	SITE	DAY 27	SITE
DAY 7	SITE	DAY 14	SITE	DAY 21	SITE	DAY 28	SITE

Still have questions about injecting Kineret or injection site reactions?

Contact KINERET® On TRACK™, read the Kineret Prescribing Information, and/or ask a healthcare professional.

CALL US: 866.547.0644, 8AM to 8PM EST, Monday through Friday.



INDICATIONS

Kineret® (anakinra) is a prescription medicine called an interleukin-1 receptor antagonist (IL-1ra) used to:

- Reduce the signs and symptoms, and slow the damage of moderate to severe active rheumatoid arthritis (RA) in people age 18 years and older when 1 or more other drugs for RA have not worked
- Treat people with a form of Cryopyrin-Associated Periodic Syndromes (CAPS) called Neonatal-Onset Multisystem Inflammatory Disease (NOMID)

IMPORTANT SAFETY INFORMATION

Who should not take Kineret?

People who are allergic to:

- proteins made from bacteria called *E. coli*. Ask your healthcare provider if you are not sure
- anakinra or any of the ingredients in Kineret. See the end of the patient leaflet for a complete list of ingredients in Kineret

What information should I know before starting Kineret?

Before you use Kineret, tell your healthcare provider if you:

- have an infection, a history of infections that keep coming back, or other problems that can increase your risk of infections
- have kidney problem
- are scheduled to receive any vaccines. People using Kineret should not receive live vaccines
- are pregnant or plan to become pregnant. It is not known if Kineret will harm your unborn baby
- are breastfeeding or plan to breastfeed. It is not known if Kineret passes into your breast milk. You and your healthcare provider should decide if you will use Kineret or breastfeed

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Kineret and other medicines may affect each other and cause serious side effects. Especially tell your healthcare provider if you take certain other medicines that affect your immune system called Tumor Necrosis Factor (TNF) Blockers. Ask your healthcare provider for a list of these medicines if you are not sure.

What are the possible side effects of Kineret?

Kineret may cause serious side effects, including:

- **serious infections.** Kineret may lower your ability to fight infections. During treatment with Kineret, call your healthcare provider right away if you get an infection, have any sign of an infection including a fever or chills, or have any open sores on your body. You may get an infection if you receive live vaccines while you use Kineret. You should not receive live vaccines while you use Kineret.
- **allergic reactions.** Stop using Kineret and call your healthcare provider or get emergency help right away if you have any of these symptoms of an allergic reaction: swelling of your face, lips, mouth, or tongue; trouble breathing; wheezing; severe itching; skin rash, redness, or swelling outside of the injections site area; dizziness or fainting; fast heartbeat or pounding in your chest (tachycardia); or sweating.

- **decreased ability of your body to fight infections (immunosuppression).** It is not known if treatment with medicines that cause immunosuppression, like Kineret, affect your risk of getting cancer.
- **low white blood cell count (neutropenia).** Kineret may cause you to have a lower number of certain white cells (neutrophils). Neutrophils are important in fighting infections. You should have blood tests before starting treatment with Kineret, then monthly for 3 months. After the first 3 months you should have your blood tested every 3 months for up to 1 year.

The most common side effects of Kineret for RA include:

- Injection site skin reactions, including redness, swelling, bruising, itching, and stinging. Most injection site reactions are mild, happen early during treatment, and last about 14 to 28 days
- rheumatoid arthritis (RA) gets worse with treatment
- sore throat or runny nose
- headache
- nausea
- diarrhea
- sinus infection
- joint pain
- feeling like you have the flu
- pain in your stomach area

The most common side effects of Kineret for NOMID include:

- Injection site skin reactions, including redness, swelling, bruising, itching, and stinging. Most injection site reactions are mild, happen early during treatment, and last about 14 to 28 days. Injection site reactions have been observed less frequently in people with NOMID
- headache
- vomiting
- diarrhea
- joint pain
- fever
- feeling like you have the flu
- sore throat or runny nose

Tell your healthcare provider if you have any side effect that bothers you or does not go away. These are not all of the possible side effects of Kineret. For more information ask your healthcare provider or pharmacist.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

